

THE THERMAL GLUEBALL AT FINITE TEMPERATURE IN SU(3) ANISOTROPIC LATTICE QCD*

NORIYOSHI ISHII

*Radiation Lab., RIKEN(The Institute of Physical and Chemical Research)
2-1 Hirosawa, Wako-shi, Saitama 351-0198, JAPAN
E-mail: ishii@rarfarp.riken.go.jp*

HIDEO SUGANUMA[†]

*Faculty of Science, Tokyo Institute of Technology,
2-12-1 Ohkayama, Meguro, Tokyo 152-8552, JAPAN*

HIDEO MATSUFURU[‡]

*Yukawa Institute for Theoretical Physics, Kyoto University,
Kitashirakawa-Oiwake, Sakyo, Kyoto 606-8502, JAPAN*

The thermal glueball is studied at finite temperature by using SU(3) anisotropic lattice QCD with $\beta_{\text{lat}} = 6.25$, the renormalized anisotropy $\xi \equiv a_s/a_t = 4$ over the lattice of the size $20^3 \times N_t$ with various N_t at the quenched level. While the narrow peak ansatz leads to the significant polemass reduction of about 300 MeV near the critical temperature T_c , Breit-Wigner ansatz which can take into account the possible appearance of the thermal width leads to the significant thermal width broadening of about 300 MeV with a modest reduction of the peak center of about 100 MeV.

At finite temperature/density, the QCD vacuum changes its structure. This is the case even in the confinement phase, where one expects the reduction in the string tension and the partial restoration of the spontaneous chiral symmetry breaking. Since hadrons are relativistic bound states of quarks and gluons, it would be natural to expect that the changes in the QCD vacuum leads to the changes in various properties of hadrons. In

*The numerical calculation has been done on NEC SX-5 at Osaka University.

[†]supported by Grant for Scientific Research (No. 12640274) from Ministry of Education, Culture, Science and Technology, Japan.

[‡]supported by Japan Society for the Promotion of Science for Young Scientists

fact, near the critical temperature T_c , a number of effective models predict the (pole)mass reductions of various hadrons of more than a few hundred MeV^{2,3,4,5}, which would serve as important precritical phenomena of the QCD phase transition. Recently, motivated by these studies, quenched anisotropic lattice QCD⁶ has been used to measure the polemass of various hadrons at finite temperature^{7,8,9}, reporting the profound results that, for both the light and heavy $q\bar{q}$ -mesons, no significant change is observed below T_c , while, for the glueball, the significant polemass reduction of about 300 MeV is observed near T_c . In all of these studies, the narrowness of the peak is assumed. However, since at $T \neq 0$, each bound state peak acquires a finite thermal width through the interaction with the heatbath, it is desirable to respect the existence of the thermal width. Here, we report an advanced analysis of the thermal 0^{++} glueball based on SU(3) quenched anisotropic lattice QCD taking into account the effect of the thermal width.

Generally, to extract physical observables such as mass and width, we have to resort to the spectral representation of the two-point correlator $G(t) \equiv Z(\beta)^{-1} \text{tr} \{ e^{-\beta H} \phi(t) \phi(0) \}$ as

$$G(t) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{\rho(\omega)}{2 \sinh(\beta\omega/2)} \cosh(\omega(\beta/2 - t)), \quad (1)$$

where H is the QCD Hamiltonian, $Z \equiv \text{tr}(e^{-\beta H})$, $\phi(t) \equiv e^{tH} \phi(0) e^{-tH}$ is the zero-momentum projected glueball operator in the imaginary-time Heisenberg picture, and $\rho(\omega)$ is the spectral function. Appropriate smearing on $\phi(t)$ is understood to maximize the overlap to the glueball state. To extract the physical observables, we parameterize $\rho(\omega)$ and use Eq. (1) to fit $G(t)$ generated by lattice QCD. In Refs. [6,7,8], the narrow-peak ansatz has been adopted, where $\rho(\omega)$ is parameterized as $\rho(\omega) = 2\pi A (\delta(\omega - m) - \delta(\omega + m)) + \dots$ with the two fit parameters A and m corresponding to the overlap and the polemass, respectively.

To respect the thermal width, we recall that $\rho(\omega)$ is the imaginary part of the retarded Green function $G_R(\omega)$, i.e., $\rho(\omega) = -2\text{Im}(G_R(\omega))$. At $T = 0$, bound state poles of $G_R(\omega)$ are located on the real ω -axis. With increasing T , they begin to move off the real axis into the complex ω -plane. Thus the influence of each complex pole in $\rho(\omega)$ amounts to a Lorentzian as $\rho(\omega) = 2\pi A (\delta_\Gamma(\omega - \omega_0) - \delta_\Gamma(\omega + \omega_0)) + \dots$, where A , ω_0 and Γ are used as fit parameters corresponding to the overlap, the center and the thermal width, respectively. $\delta_\epsilon(x) \equiv \frac{1}{\pi} \text{Im} \left(\frac{1}{x - i\epsilon} \right)$ denotes a smeared delta function. We refer to the corresponding fit function as the Breit-Wigner type.

The numerical results are shown in Fig. 1. We use 5000 to 9900 gauge

configurations generated by the Wilson action with $\beta = 6.25$ and the renormalized anisotropy $\xi \equiv a_s/a_t = 4$. Whereas the narrow-peak ansatz indicates the significant polemass reduction of about 300 MeV, the Breit-Wigner ansatz indicates the significant thermal width broadening of more than 300 MeV with a modest reduction in the peak center. These two analyses thus lead to the two different physical implications. Note that, due to the biased factor “ $\sinh(\beta\omega/2)$ ” in Eq. (1) which enhances the smaller ω region of $\rho(\omega)$, thermal width is effectively seen as the reduced polemass in the narrow-peak ansatz. Hence, in the case of the glueball, the thermal effect is most probably the thermal width broadening.




Figure 1. The polemass $m_G(T)$ from the narrow-peak ansatz, the center $\omega_0(T)$ and the thermal width $\Gamma(T)$ from the Breit-Wigner ansatz. The vertical dotted lines indicate the critical temperature $T_c \simeq 280$ MeV. Appropriate smearings are understood.

To summarize, we have studied the thermal glueball by using anisotropic lattice QCD at quenched level. While the narrow-peak ansatz has lead to the significant polemass reduction near T_c , the Breit-Wigner analysis has lead to the significant thermal width broadening with a modest reduction in the peak center. Since the thermal width can be effectively seen as the polemass reduction, we have concluded that the thermal effect on the glueball is most probably the thermal width broadening.

References

1. N. Ishii et al., Phys. Rev. **D66**, 094506 (2002).
2. T. Hatsuda et al., Phys. Rev. Lett. **55**, 158 (1985).
3. T. Hashimoto et al., Phys. Rev. Lett. **57**, 2123 (1986).
4. T. Hatsuda et al., Phys. Rev. **D47**, 1225 (1993).
5. H. Ichie et al., Phys. Rev. **D52**, 2944 (1995).
6. T. R. Klassen, Nucl. Phys. **B533**, 557 (1998).
7. QCD-TARO Collaboration, Ph. de Forcrand, Phys. Rev. **D63**, 054501 (2001).
8. T. Umeda et al., Int. J. Mod. Phys. **A16**, 2215 (2001).
9. N. Ishii et al., Phys. Rev. **D66**, 014507 (2002).